

Chapter Thirteen

DISEASES OF INTERIORSCAPE PLANTS

GENERAL METHODS OF DISEASE CONTROL	3
CACTUS (Family Cactaceae)	6
Gray Mold	6
Leaf Spot And Stem Rot.....	6
CHINESE EVERGREEN (<i>Algaonema</i> spp.)	7
BACTERIAL LEAF SPOT	7
CROTON (<i>Cordiaecum variegatum</i>)	7
Anthracnose	7
DIEFFENBACHIA (<i>Dieffenbachia</i> spp.).....	7
Erwinia Blight And Stem Rot.....	8
Leaf Scorch.....	8
Root And Stem Rot	9
Xanthomonas Leaf Spot	9
DRACAENA (<i>Dracaena</i> spp.)	9
Botrytis Blight.....	9
Fusarium Leaf Spot And Stem Rot.....	10
Leaf Scorch.....	11
ENGLISH IVY (<i>Hedera helix</i>)	11
Bacterial Leaf Spot.....	11
FERN (Family Polypodiaceae)	11
Bacterial Leaf Blight.....	11
FIG (<i>Ficus</i> spp.).....	12
Phomopsis Dieback.....	12
GRAPE IVY, KANGAROO VINE (<i>Cissus</i> spp.)	12
Powdery Mildew.....	12
Root Rot.....	13
INDIA RUBBER TREE (<i>Ficus elastica</i>).....	13
Anthracnose	13
Leaf Scorch.....	13
NORFOLK ISLAND PINE (<i>Araucaria heterophylla</i>)	14
Colletotrichum Needle Blight	14

Gray Mold	14
PALM (family Palmaceae)	14
Fusarium Wilt	14
Gray Mold	15
Leaf Scorch	15
PEACE LILY (<i>Spathiphyllum</i> spp.)	15
Cylindrocladium Root And Petiole Rot	15
PEPEROMIA (<i>Peperomia obtusifolia</i>)	16
Leaf Scorch	16
Stem And Root Rot	16
PHILODENDRON (<i>Philodendron</i> spp.)	17
Erwinia Blight	17
Leaf Scorch	17
Pseudomonas Leaf Spot	17
Stem And Root Rot	18
SCHEFFLERA (<i>Brassaia actinophylla</i>)	18
Leaf Scorch	18
Root-Knot Nematode	18
GENERAL REFERENCES	20
APPENDIX I. Fungicides registered for use against diseases of foliage plants in Canada	22
APPENDIX II. Latin and common names of foliage plants	28

GENERAL METHODS OF DISEASE CONTROL

Introduction

One of the fastest-growing segments of modern-day commercial horticulture is ornamental plant production. Within this specialized industry, the use of living plants in interior locations such as homes, apartments, offices, public buildings and shopping malls has rapidly expanded in recent years. Many of these plants are of tropical origin, so the interior environments in which they are placed are markedly different than their native habitats. This may place the plants under stress and thereby predispose them to a variety of infectious and non-infectious diseases. Preventing and remedying these problems can be a challenging task because of the difficulty of applying conventional disease control practices in the interior landscape (interiorscape).

Influence of Production Practices on Diseases

Foliage plants are produced under a wide range of environmental and cultural conditions. In the southern United States, many species are grown outdoors in full sun or under varying degrees of artificial shade. In northern areas of the U.S. and Canada, enclosed structures such as greenhouses are widely used. Many of the same diseases prevail throughout the major production areas. Likewise, common problems are encountered in interiorscapes, regardless of their geographical location. Major changes in production practices have decreased the importance of a number of infectious diseases. For example, the increased use of sterilized growing media and containers has prevented many root diseases from becoming epidemic. The availability of effective, modern pesticides has been of great help to foliage plant producers, as has access to pathogen-free, tissue-cultured plantlets.

The far-ranging and rapid transport of foliage plants throughout the world poses some unique problems in controlling diseases. Pathogens may be introduced into plantings in a variety of ways. Cuttings and other propagative parts are often produced in the tropics and transported to new production locations, complete with diseases and pests. In other cases, propagation units may be derived from infected stock plants produced by a grower for his own use or for sale to other growers. Further spread of diseases may occur during retail distribution. Some tropical plant diseases cannot develop in temperate areas because environmental conditions are not suitable. However, other diseases may flourish over a wide range of conditions, especially if the plants are produced in warm and/or humid environments. Control strategies, therefore, may differ tremendously depending upon the stage at which plants become infected and develop symptoms, and the regime under which they are being grown. Rigorous and routine examination of plant material at all stages of production and utilization should enable prompt and accurate problem diagnoses and the development of effective disease management strategies.

Cultural Control Measures

Major steps towards effective disease control can be taken with simple changes in existing cultural methods. Some of these are listed below and others are discussed in Chapter 4 under 'General Disease Control Methods for Greenhouse Crops'.

- a) Employ watering systems that do not wet the foliage. Many fungal and bacterial pathogens depend upon free moisture for spore dispersal, germination and/or infection.
- b) Water as infrequently as possible, only when needed, and thoroughly enough to wet the root ball. Extra water may be required to leach out excess soluble salts in some situations.
- c) Water when good drying conditions exist, i.e. not too early or late in the day.

- d) Employ fans, where needed, to move drier air in and around plants to promote foliar drying.
- e) Avoid overcrowding plants. This promotes poor drying and extends wet periods by increasing humidity around plants, thereby causing condensation to occur on plant surfaces. Crowding may also promote tissue senescence, which predisposes it to attack by certain fungi, e.g. *Botrytis*.
- f) Use water free of bacterial and fungal pathogens and low in soluble salts and other impurities.
- g) Pasteurize media containing soil, sand and/or peat moss using heat or chemicals. Holding the temperature at 82°C (180°F) for at least 30 minutes will kill most pathogens.
- h) Clean and disinfest pots, tools, hands and other items coming in contact with pathogen-free media and plants. Benches used for propagation and/or potted plants should be washed and steam or chemically treated prior to receiving media or pots. It is also desirable to raise the benches above the ground to avoid possible contamination from the soil below.
- i) Propagate and install plant stocks which are known to be pathogen-free. Badly infected plants should be rogued and destroyed.
- j) Use growing media with adequate pore space to facilitate good aeration and rapid drainage. Provide sufficient subsurface drainage to allow for leaching out of excess soluble salts.
- k) Provide adequate levels of essential nutrients for plant growth under the particular growing regimes in use. Avoid over-fertilizing as this promotes the accumulation of soluble salts in growing media. Adjustments in nutrient levels so as to promote slower, harder growth will decrease the risk of bacterial diseases that generally thrive on soft, succulent foliage.
- l) Provide optimum air and soil temperatures and light levels for the particular species being grown. Low temperatures and inadequate light result in weak, spindly growth and may predispose plants to attack by pathogens by reducing their natural resistance.
- m) Remove infected leaves and branches, where practical. Gather and destroy prunings and fallen leaves and branches.
- n) Toxic levels of fluorides can occur in superphosphate fertilizer, perlite, water and some peats. Addition of dolomitic or calcitic lime to potting media to increase the pH reduces the solubility of fluoride. Sensitive plants should be grown in a medium of pH 6.0 or greater.

Chemical Control Measures

The requirement for effective disease control chemicals is greatest in the propagation and growing-on phases of foliage plant production. Here, conditions are more favorable for disease development and spread than in interiorscapes. At present in Canada, there are no fungicides, bactericides or soil fumigants specifically registered for use in interiorscapes. Those products which are licensed on foliage plants are only to be used in greenhouses or, in a few situations, households. Growers should evaluate their individual situations and determine chemical control needs based on potential or actual disease problems. As well, they should guard against overprotection and misuse of pesticides as such practices may be costly, hazardous to human health, and potentially harmful to the plants themselves.

- a) Foliar Fungicides - The number of fungal pathogens that can attack the tops of foliar plants is extensive. Preventative, bi-weekly sprays to both leaf surfaces with broad-spectrum fungicides should provide adequate protection against fungal attack. Foliar sprays may leave some residue on the foliage, especially if spreader-stickers are used. A list of the fungicides registered specifically on foliage plants in Canada is given in Appendix I.
- b) Foliar Bactericides - At present, there are no pesticides registered in Canada specifically the control of bacterial diseases on foliage plants.
- c) Soil Fungicides - The largest amount of fungicide use associated with foliage plant production occurs in the control of soil-borne plant pathogens. Growers face the prospect of controlling many different soil-borne plant pathogens on a wide variety of hosts. Soil fungicides are generally applied as drenches. The volume of drench used depends largely upon the size of container, depth of bed and type of growing media, especially the amount of organic matter. There are often difficulties encountered when applying soil fungicides that may result in poor control or phytotoxicity. Biodegradable, nonionic surfactants may aid in fungicide distribution and disease control. Generally, drenches should not be used more frequently than once every three months. Also, they should not be applied to the foliage. If a misapplication occurs, lightly rinse the foliage with water to remove the material. Where drenches are used in cutting beds, they should be applied prior to sticking the cuttings. Biofungicides are also now available for use in greenhouse crops to help suppress root rotting organisms such as *Pythium*, *Rhizoctonia* and *Fusarium*.
- d) Nematicides - There are no nematicides registered in Canada for nematode control specifically on foliage plants.

Resistant Cultivars

The relative resistance of various species of foliage plants to prevalent infectious and non-infectious diseases has not been well documented. Foliage plant breeders have placed relatively little emphasis on incorporating disease resistance into newly developed cultivars. Likewise, plant pathologists have done only limited evaluations of the relative resistance of foliage plant cultivars currently in commercial use. Until more work is done, the use of resistant cultivars for disease management in foliage plants will not be a readily available alternative for foliage plant producers and interior landscapers.

CACTUS (Family Cactaceae)

GRAY MOLD

Botrytis cinerea

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Fenhexamid (COM)WDG. See Appendix I. Limitations: As per label.

LEAF SPOT AND STEM ROT

Fusarium oxysporum, Drechslera cactivora, Pythium aphanidermatum, Phytophthora parasitica

Cultural: Rogue infected plants. Avoid mechanical injury. Do not over water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: See table 1.

Table 1. Relative susceptibility of holiday cacti to four plant pathogens (1).¹

Cultivar	<i>Erwinia</i>	<i>Drechslera</i>	<i>Fusarium</i>	<i>Phytophthora</i>
Bridgeport	moderate	moderate	low	moderate
Cambridge	high	moderate	variable	high
Christmas Charm	variable	moderate	moderate	moderate
Christmas Fantasy	low to moderate	moderate	variable	variable
Christmas Flame	variable	moderate	high	moderate
Christmas Magic II	low	moderate	low	low
Gold Charm	high	variable	high	high
Holiday Splendor	variable	moderate	low	high
Kris Kringle	low	high	moderate	variable
Kris Kringle II	low to moderate	high	variable	variable
Lavender Doll	variable	low to moderate	low	variable
Lavender Doll II	low	variable	variable	moderate to high
Peach Parfait	variable	high	low	high
Red Radiance	variable	moderate	moderate	variable
Sanibel	variable	variable	low	moderate
Santa Cruz	high	low to moderate	moderate	moderate
Sleigh Bells	variable	moderate	low	variable
Twilight Tangerine	high	moderate	moderate	moderate
White Christmas	low	low to moderate	moderate	moderate
Windsor	variable	moderate	variable	variable

¹ *Erwinia carotovora* subsp. *carotovora*, *Drechslera cactivora*, *Fusarium oxysporum*, *Phytophthora parasitica*

Chemical: None.

References:

1. Chase, A.R. 1982. Stem rot and shattering of Easter cactus caused by *Drechslera cactivora*. Plant Dis. 66: 602-603.
2. Chase, A.R. 1993. Susceptibility of holiday cactus cultivars to diseases. Foliage Digest 14(11): 7-8.
3. Knauss, J.F. 1975. Control of basal stem and root rot of Christmas cactus caused by *Pythium aphanidermatum* and *Phytophthora parasitica*. Proc. Fla. State Hort. Soc. 88: 567-571.
4. Mitchell, J.K. 1987. Control of basal stem and root rot of Christmas and Easter cacti caused by *Fusarium oxysporum*. Plant Dis. 71: 1018-1020.

CHINESE EVERGREEN (*Algaonema* spp.)

BACTERIAL LEAF SPOT

Erwinia chrysanthemi, *Pseudomonas cichorii*, *Xanthomonas campestris* pv. *dieffenbachiae*

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Arias, R.S. *et al.* 1998. Rapid detection of pectolytic *Erwinia* spp. in *Aglaonema* sp. HortTechnology 8(4):602-605.
2. Chase, A.R. and Poole, R.T. 1986. Troubleshooting guide to foliage: aglaonema. Greenhouse Grower 4(4):54-55.
3. McFadden, L.A. 1962. Two bacterial pathogens affecting leaves of *Aglaonema robelinii*. Phytopathology 52:20. (Abstr.)
4. McFadden, L.A. 1969. *Aglaonema pictum*, a new host for *Erwinia chrysanthemi*. Plant Dis. Rep. 53: 253-254.

CROTON (*Cordiaum variegatum*)

ANTHRACNOSE

Glomerella cingulata

Cultural: Prune off infected leaves or cut away infected areas of leaves. Disinfect pruning tools after use to avoid spreading the pathogen to healthy tissue. Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Osborne, L.S. *et al.* 1984. Croton. Foliage Digest 7(4): 5-7.

DIEFFENBACHIA (*Dieffenbachia* spp.)

ERWINIA BLIGHT AND STEM ROT

Erwinia carotovora subsp. *carotovora*, *E. chrysanthemi*

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Henny, R.J., A.R. Chase and L.S. Osborne. 1994. *Dieffenbachia*. Foliage Digest 17(6): 3-6.
2. McFadden, L.A. 1961. Bacterial stem and leaf rot of *Dieffenbachia* in Florida. Phytopathology 51: 663-668.
3. Munnecke, D.E. 1960. Bacterial stem rot of *Dieffenbachia*. Phytopathology 50: 698-700.
4. Nieves-Brun, C. 1985. Infection of roots of *Dieffenbachia maculata* by the foliar blight and soft rot pathogen, *Erwinia chrysanthemi*. Plant Pathol. 34: 139-145.

LEAF SCORCH

Physiological - Excess soluble salts, overwatering, poor drainage, excess boron or fluorides.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (1). Carefully regulate fertilizer applications. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Henny, R.J., A.R. Chase and L.S. Osborne. 1994. *Dieffenbachia*. Foliage Digest 17(6): 3-6.
2. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton, AB.
3. Poole, R.T. and Chase, A.R. 1987. *Dieffenbachia* guide. Greenhouse Grower 5(3): 32-34.

ROOT AND STEM ROT

Phytophthora spp., *Pythium* spp., *Fusarium solani*

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Etridiazole (COM) WP, EC for *Pythium* and *Phytophthora*. *Trichoderma harzianum* (COM) GR, WP for *Pythium*, *Rhizoctonia*, *Fusarium*. *Streptomyces griseoviridis* (COM) WP for *Fusarium* diseases. See Appendix I. Limitations: As per label.

References:

1. Chase, A.R. and El-Gholl, N.E. 1982. Stem rot, cutting rot, and leaf spot of *Dieffenbachia maculate* 'Perfection' incited by *Fusarium solani*. Plant Dis. 66: 595-598.
2. Chase, A.R. 1985. You can prevent foliage diseases. Greenhouse Grower 3(12): 38-45.
3. Henny, R.J., A.R. Chase and L.S. Osborne. 1994. Dieffenbachia. Foliage Digest 17(6): 3-6.

XANTHOMONAS LEAF SPOT

Xanthomonas campestris pv. *dieffenbachiae*

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: Tropic Star.

Intermediate: Bausei, Camille, Compacta, Memoria-Corsii, Star White, Starry Nights.

Susceptible: Triumph, Victory

Chemical: None.

References:

1. Chase, A.R. 1985. You can prevent foliage diseases. Greenhouse Grower 3(12): 38-45.
2. Chase, A.R. and Henny, R.J. 1990. Susceptibility of 12 *Dieffenbachia* cultivars to *Xanthomonas campestris* pv. *dieffenbachiae*. Foliage Digest 13(2): 1-2.
3. Henny, R.J., A.R. Chase and L.S. Osborne. 1994. Dieffenbachia. Foliage Digest 17(6): 3-6.

DRACAENA (*Dracaena* spp.)

BOTRYTIS BLIGHT

Botrytis cinerea

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Fenhexamid (COM) WDG; iprodione (COM) WP. See Appendix I. Limitations: As per label.

FUSARIUM LEAF SPOT AND STEM ROT

Fusarium moniliforme

Cultural: Keep the foliage as dry as possible to reduce the risk of infecting new leaves. The severity of Fusarium leaf spot on red-edge dracaena (*Dracaena marginata*) can be substantially reduced by using increased rates of fertilizer

(1). Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: Resistance of selected *Dracaena* species and cultivars to Fusarium leaf spot (1).

<i>Dracaena</i> species and cultivars	Resistant or Susceptible	Symptom type
<i>deremensis</i> 'Compacta'	resistant	none
<i>deremensis</i> 'Janet Craig'	very slightly susceptible	clear, yellow speckles on leaf edges
<i>deremensis</i> 'Warneckii'	very slightly susceptible	clear, yellow speckles on leaf edges
<i>deremensis</i> 'Lemon Lime'	resistant	none
<i>fragrans</i> 'Massangeana'	very slightly susceptible	small yellow spots on leaf edges
<i>marginata</i>	very highly susceptible	large yellow and brown spots in whorl that can blend to rot center
<i>marginata</i> 'Bicolor'	very slightly susceptible	small yellow spots in whorl
<i>marginata</i> 'Colorama'	moderately susceptible	small yellow spots in whorl
<i>marginata</i> 'Magenta'	very highly susceptible	large brown spots in whorl that can blend to rot center
<i>marginata</i> 'Tricolor'	slightly susceptible	small yellow spots in whorl
<i>reflexa</i> 'Song of Jamaica'	highly susceptible	large tan to reddish brown spots in whorl and on leaf edges
<i>sanderana</i>	resistant	none
<i>sanderana</i> 'Borinquensis'	resistant	none
<i>sanderana</i> 'Gold'	resistant	none
<i>surculosa</i>	very slightly susceptible	large tan papery spots form where leaf joins stem
<i>surculosa</i> 'Florida Beauty'	very slightly susceptible	large tan papery spots form where leaf joins stem
<i>surculosa</i> 'Juanita'	very slightly susceptible	large tan papery spots form where leaf joins stem

Chemical: Chlorothalonil (COM) SU, WDG. See Appendix I. Limitations: As per label.

References:

1. Chase, A.R. 1987. Effect of fertilizer level on severity of Fusarium leaf spot of *Dracaena marginata*. Proc. Fla. State Hort. Soc. 100: 360-362.
2. Chase, A.R. 1993. Fusarium leaf spot of Dracaenas: resistance of species and cultivars. Foliage Digest 16(8): 1-2.
3. Chase, A.R. 1990. Fertilizer rate affects severity of Fusarium leaf spot of red-edge dracaena. Foliage Digest 13(6): 6-7.
4. Knauss, J.F. 1971. Fusarium stem rot, a previously unreported disease of *Dracaena*. Proc. Am. Soc. Hort. Sci., Trop. Reg. 15: 208-215.
5. Poole, R.T., A.R. Chase and L.S. Osborne. 1993. Dracaena. Foliage Digest 16(11): 1-6.
6. Wehlburg, C. and Martinez, A.P. 1967. Leaf spot of *Dracaena marginata* Lam. caused by *Fusarium moniliforme* Sheld. Proc. Fla. State Hort. Soc. 80: 454-456.

LEAF SCORCH

Physiological - Excess soluble salts, overwatering, poor drainage, excess boron or fluorides.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (2). Carefully regulate fertilizer applications. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Conover, C.A. and Poole, R.T. 1982. Fluoride induced chlorosis and necrosis of *Dracaena fragrans* 'Massangeana'. J. Amer. Soc. Hort. Sci. 107: 136-139.
2. Conover, C.A. and Poole, R.T. 1980. Influence of fertilization, superphosphate and lime on mottling of *Dracaena fragrans*. HortScience 15: 23-24.
3. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton,

ENGLISH IVY (*Hedera helix*)

BACTERIAL LEAF SPOT

Xanthomonas campsetris pv. *hederae*

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: Gold Dust, Perfection, Sweet Heart, Eva, California.

Intermediate: Telecurl, Manda Crested, Gold Heart.

Susceptible: Ivalace, Green Variegated, Hahn Variegated, Brokamp.

Chemical: None.

References:

1. Chase, A.R. 1984. *Xanthomonas campestris* pv. *hederae* causes a leaf spot of five species of Araliaceae. Plant Pathol. 33: 439-440.
2. Norman, D.J. 1996. Diseases of English ivy. Foliage Digest 19(5): 1-2.
3. Osborne, L.S. and Chase, A.R. 1985. Susceptibility of cultivars of English ivy to two-spotted spider mite and *Xanthomonas* leaf spot. HortScience 20: 269-271.

FERN (Family Polypodiaceae)

BACTERIAL LEAF BLIGHT

Pseudomonas cichorii, *P. gladioli*

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Ark, P.A. and Thomkins, C.M. 1946. Bacterial leaf blight of bird's-nest fern. *Phytopathology* 36: 758-761.
2. Chase, A.R. *et al.* 1984. Leaf spot and blight of *Asplenium nidus* caused by *Pseudomonas gladioli*. *Plant Dis.* 68: 344-347.

FIG (*Ficus* spp.)

PHOMOPSIS DIEBACK

Phomopsis cinerescens

Cultural: Plant healthy trees and follow good growing practices to minimize stress. Prune dead or dying branches, disinfecting shears between cuts. Destroy all leaf and twig litter. Maintain light levels at 150 ft-c or more. Follow natural target pruning guidelines in order to favour compartmentalization and inhibit *Phomopsis* infection.

Resistant Cultivars: None. See Notes.

Chemical: None.

Notes: *Ficus benjamina* (weeping fig) is particularly susceptible to phomopsis dieback, whereas other *Ficus* spp. appear less susceptible (3).

References:

1. Anderson, R.G. and Hartman, J.R. 1983. Phomopsis twig blight on weeping fig indoors: a case study. *Foliage Digest* 6: 5-7.
2. Benschop, K., Tewari, J.P. and Toop, E.W. 1984. Phomopsis twig dieback of some woody interior ornamentals in Alberta. *Can. Plant Dis. Surv.* 64: 29-31.
3. Chase, A.R. 1993. Common diseases of ficus. *Southern Nursery Digest* (February issue): 20-21.
4. Henley, R.W., A.R. Chase and L.S. Osborne. 1994. Ficus. *Foliage Digest* 17(2): 1-6.
5. Newman, L. 1991. The ultimate *Ficus* prune. *Interiorscape* 10(1): 6-7.
6. Newman, L. 1993. Ficus pruning in the interiorscape. *Foliage Digest* 14(10): 7-8.

GRAPE IVY, KANGAROO VINE (*Cissus* spp.)

POWDERY MILDEW

Oidium spp.

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Henley, R.W., A.R. Chase and L.S. Osborne. 1993. Grape ivy. Foliage Digest 14(10): 1-4.

ROOT ROT

Pythium spp., *Rhizoctonia solani*, *Fusarium oxysporum*, *Cylindrocarpon* spp.

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: *Trichoderma harzianum* (COM) GR, WP for *Pythium*, *Rhizoctonia*, and *Fusarium*; *Streptomyces griseoviridis* (COM) WP for *Fusarium* diseases. See Appendix I. Limitations: As per label.

References:

1. Bolton, A.T. 1984. Root rot of *Ficus benjamina*. Plant Disease 68: 816-817.

INDIA RUBBER TREE (*Ficus elastica*)

ANTHRACNOSE

Glomerella cingulata

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Agrawal, S.C. and Saksena, S.B. 1972. Experimental studies on leaf anthracnose of *Ficus elastica* Roxb. Curr Sci. 41: 246-249.
2. Henley, R.W., A.R. Chase and L.S. Osborne. 1994. Ficus. Foliage Digest 17(2): 1-6.

LEAF SCORCH

Physiological - Excess soluble salts, overwatering, poor drainage, excess boron or fluorides.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (1). Carefully regulate fertilizer application. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Henley, R.W., A.R. Chase and L.S. Osborne. 1994. *Ficus*. Foliage Digest 17(2): 1-6.
2. Marlatt, R.B. 1978. Boron deficiency and toxicity symptoms in *Ficus elastica* 'Decora' and *Chrysalidocarpus lutescens*. HortScience 13: 442-443.
3. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton, AB.

NORFOLK ISLAND PINE (*Araucaria heterophylla*)

COLLETOTRICHUM NEEDLE BLIGHT

Colletotrichum derridis

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Ridings, W.H. 1973. Colletotrichum needle necrosis of Norfolk Island Pine. Proc. Fla. State Hort. Soc. 86: 418-421.

GRAY MOLD

Botrytis cinerea

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Fenhexamid (COM) WDG. See Appendix I. Limitations: As per label.

PALM (family Palmaceae)

FUSARIUM WILT

Fusarium oxysporum

Cultural: Disinfest pruning tools between plants. Also refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: *Streptomyces griseovirdis* (COM) WP. See Appendix I. Limitations: As per label.

References:

1. Chase, A.R. 1993. Common diseases of palms. Southern Nursery Digest (March issue): 20-21.
2. Domner, J. and Ohr, H. 1989. Palm trees have problems, too. Grounds Maintenance 24(4): 94-102.

GRAY MOLD

Botrytis cinerea

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Chlorothalonil (COM) PO; fenhexamid (COM) WDG. See Appendix I. Limitations: As per label.

Notes: Apply chlorothalonil only when the foliage is dry, and when air temperatures do not exceed 24°C.

LEAF SCORCH

Physiological - Excess soil salts, overwatering, poor drainage, excess fluorides or boron.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (1). Carefully regulate fertilizer application. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Chase, A.R. and Poole, R.T. 1986. Troubleshooting guide to foliage: palms. Greenhouse Grower 4(11):24-27.
2. Henley, R.W., A.R. Chase and L.S. Osborne. 1993. Chamaedorea palm. Foliage Digest 16(9): 1-4.
3. Marlatt, R.B. 1978. Boron deficiency and toxicity symptoms in *Ficus elastica* 'Decora' and *Chrysalidocarpus lutescens*. HortScience 13: 442-443.
4. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton, AB.

PEACE LILY (*Spathiphyllum* spp.)

CYLINDROCLADIUM ROOT AND PETIOLE ROT

Cylindrocladium spathiphylli

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Chase, A.R. and Poole, R.T. 1988. *Cylindrocladium* root and petiole rot of *Spathiphyllum* spp. Bull. 860, Agric. Exp. Sta., Univ. Fla., Gainesville, FL. 20 pp.
2. Schoulties, C.L. and El-Gholl, N.E. 1983. Host range and pathogen specificity studies of *Cylindrocladium spathiphylii*. Proc. Fla. State Hort. Soc. 96: 282-284.

PEPEROMIA (*Peperomia obtusifolia*)

LEAF SCORCH

Physiological - Excess soluble salts, overwatering, poor drainage, excess boron or fluoride.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (1). Carefully regulate fertilizer application. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton, AB.
2. Poole, R.T. and Chase, A.R. 1987. Response of foliage plants to fertilizer application rates and associated leachate conductivity. HortScience 22: 317-318.

STEM AND ROOT ROT

Rhizoctonia solani, *Phytophthora* spp., *Pythium* spp.

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Etridiazole (COM) WP, EC; *Trichoderma harzianum* (COM) GR, WP. See Appendix I.
Limitations: As per label.

Notes: Etridiazole will only control *Pythium* and *Phytophthora*; *Trichoderma harzianum* will control *Pythium*, *Rhizoctonia* and *Fusarium*.

References:

1. Ark, P.A. and DeWolfe, T.A. 1951. *Phytophthora* rot of *Peperomia*. Plant Dis. Rep. 60: 632-633.
2. Chase, A.R. and Munnecke, D.E. 1978. *Pythium* root rot of *Peperomia obtusifolia* var. *variegata*. Plant Dis. Rep. 62: 314-315.

3. Chase, A.R. and Poole, R.T. 1984. Investigations into the role of fertilizer level and irrigation frequency on growth, quality, and severity of *Pythium* root rot of *Peperomia obtusifolia*. J. Amer. Soc. Hort. Sci. 109: 619-622.

PHILODENDRON (*Philodendron* spp.)

ERWINIA BLIGHT

Erwinia chrysanthemi

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Chase, A.R. 1985. You can prevent foliage diseases. Greenhouse Grower 3(12): 38-45.
2. Chase, A.R. 1993. Common diseases of philodendron. Southern Nursery Digest (May issue): 20-21.
3. Henley, R.W., A.R. Chase and L.S. Osborne. 1994. Philodendrons: self-heading types. Foliage Digest 17(9): 4-8.
4. Knauss, J.F. and Miller, J.W. 1974. Etiological aspects of bacterial blight of *Philodendron selloum* caused by *Erwinia chrysanthemi*. Phytopathology 64: 1526-1528.

LEAF SCORCH

Physiological - Excess soluble salts, overwatering, poor drainage, excess fluorides or boron.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (1). Carefully regulate fertilizer application. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton, AB.
2. Poole, R.T. and Chase, A.R. 1987. Heartleaf philodendron guide. Greenhouse Grower 3(12): 38-45.

PSEUDOMONAS LEAF SPOT

Pseudomonas cichorii

Cultural: Avoid misting or syringing leaves with water. Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

References:

1. Chase, A.R. 1985. You can prevent foliage diseases. *Greenhouse Grower* 3(12): 38-45.
2. Henley, R.W., A.R. Chase and L.S. Osborne. 1994. Philodendrons: self-heading types. *Foliage Digest* 17(9): 4-8.

STEM AND ROOT ROT

Phytophthora spp.

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: Etridiazole (COM) WP, EC. See Appendix I. Limitations: As per label.

References:

1. Chase, A.R. 1985. You can prevent foliage diseases. *Greenhouse Grower* 3(12): 38-45.

SCHEFFLERA (*Brassaia actinophylla*)

LEAF SCORCH

Physiological - Excess soil salts, overwatering, poor drainage, excess fluorides or boron.

Cultural: Use demineralized water for irrigation and leaching. Repot when accumulations of salt are noticed on the soil surface or when scorch symptoms appear. Avoid overwatering by allowing the soil to dry between applications. In areas of poor drainage, apply sugar solution (20 g sucrose/L water) at a rate of 9 L/m² bed area (1). Carefully regulate fertilizer application. Avoid adding fluoride-containing amendments to growing media.

Resistant Cultivars: None.

Chemical: None.

References:

1. Chase, A.R. and Poole, R.T. 1986. Troubleshooting guide to foliage: schefflera. *Greenhouse Grower* 4(10): 68-69.
2. Mirza, M. 1999. Personal communication. Crop Diversification Centre North, Edmonton, AB.
3. Poole, R.T. and Chase, A.R. 1987. Response of foliage plants to fertilizer application rates and associated leachate conductivity. *HortScience* 22: 317-318.
4. Poole, R.T., A.R. Chase and L.S. Osborne. 1993. Schefflera. *Foliage Digest* 14(12): 4-8.

ROOT-KNOT NEMATODE

Meloidogyne spp.

Cultural: Refer to 'General Methods of Disease Control' at the beginning of this chapter.

Resistant Cultivars: None.

Chemical: None.

GENERAL REFERENCES

1. Alfieri, S.A., Jr., Langdon, K.R., Wehelburg, C., and Kimbrough, J.W. 1984. Index of plant diseases in Florida. Florida Dept. of Agric. and Consumer Serv., Bull. 11. 389 pp.
2. Calvert, R. (ed.). 1987. Florida foliage locator: 1987-88. Fla. Foliage Assoc., Apopka, FL. 188 pp.
3. Carroll, J. and Daughtrey, M. 1986. Diagnosis and management of disease and foliage plants in interiorscapes. *Plant Diagnostician's Quarterly*, Vol. VII, No. 2, June.
4. Chase, A.R. 1987. Compendium of ornamental foliage plant diseases. APS Press. The American Phytopathological Society. 92 pp.
5. Chase, A.R. 1991. Rhizoctonia diseases of Florida ornamentals. *Foliage Digest* 14(6): 1-3.
6. Chase, A.R. 1992. Using temperature information to aid in controlling diseases of ornamentals. *Foliage Digest* 15(2): 5-7.
7. Chase, A.R. 1993. Indoor plant problems. Part I. *Florida Growers' Ornamental Outlook* 2(6): 16.
8. Chase, A.R. 1993. Indoor plant problems. Part II. *Florida Growers' Ornamental Outlook* 2(7): 16-17.
9. Chase, A.R. 1997. Foliage plant diseases: diagnosis and control. APS Press. The American Phytopathological Society, St. Paul, Minn. 180 pp.
10. Chase, A.R. and Broschat, T.K. (eds.). 1991. Diseases and disorders of ornamental palms. APS Press, St. Paul, Minn. 56 pp.
11. Chase, A.R. and Conover, C.A. 1992. Using temperature information to aid in controlling diseases of ornamentals. *Foliage Digest* 15(2): 5-7.
12. Conner, D. and Jones, H.L. 1982. Plant pathology circular (1-200) Set 1. Bureau of Plant Pathology, Florida Dept. of Agriculture and Consumer Services, Division of Plant Industry.
13. Engelhard, A.W. 1991. Using IPM for disease control. *Foliage Digest* 14(2): 1-5.
14. Forsberg, J.L. 1975. Diseases of ornamental plants. Univ. Ill. Spec. Publ. 3 (rev.). 220 pp.
15. Gerbus, M.E., Stone, A.G. and Hoitink, A.J. 1996. Control of diseases caused by soilborne plant pathogens. *Foliage Digest* 19(3): 4-7.
16. Henley, R.W. 1983. A pictorial atlas of foliage plant problems. Central Chapter, Florida Foliage Association, Apopka, FL. 40 pp.
17. Howard, R.J. and Dykstra, M.D. 1987. Diseases of interiorscape plants in Alberta. *Agriculture and Forestry Bull.* 10(1): 22-24.
18. Joiner, J.N. 1981. Foliage plant production. Prentice-Hall, Inc., Englewood Cliffs, N.J. 614 pp.
19. Lindquist, R.K. and C.C. Powell. 1991. Pesticide application techniques for disease and insect management. *Foliage Digest* 14(12): 4-6.

20. Marlatt, R.B. 1980. Noncontagious diseases of tropical foliage plants. Agricultural Experiment Station. Institute of Food and Agric. Sciences, Univ. of Florida, Gainesville. Bull. 812.
21. Newman, L. 1996. What's killing our trees. *Interiorscape* 15(5): 20-29.
22. Newman, L. 1997. Avoid winter worries by planning holiday plant care. *Interiorscape* 16(4): 24-28.
23. O'Bannon, J.H., *et al.* 1992. Guidelines for compost sanitation. *Foliage Digest* 15(4): 7-8.
24. Pirone, P.P. 1978. Diseases and pests of ornamental plants (5th ed.). John Wiley and Sons, New York, N.Y. 566 pp.
25. Powell, C.C. 1991. Those "other" leaf spots, blights and cankers. *Foliage Digest* 14(4): 3-6.
26. Powell, C.C. 1992. Botrytis blights of flowers and foliage plants in the greenhouse. *Foliage Digest* 15(7): 6-7.
27. Powell, C.C. 1994. Infectious diseases of plants in interiorscapes. Part I: powdery mildew. *Foliage Digest* 17(8): 6-8.
28. Powell, C.C. and Lindquist, R.K. 1984. Pest and disease control on indoor plants. Cooperative Extension Service, The Ohio State University. Bull. 711, Agdex 280/605.
29. Strider, D.L. 1985. Diseases of floral crops. Vols. 1 & 2. Praeger Publ., New York, N.Y.
30. Thompson, W.T. 1994. Agricultural chemicals. Book III - Fumigants, Growth Regulators, Repellants, Rodenticides (1994-95 Revision). Thompson Publications, Fresno, Calif. 210 pp.
31. Thompson, W.T. 1997. Agricultural chemicals. Book IV. Fungicides (1997 Revision). Thompson Publications, Fresno, Calif. 235 pp.

APPENDIX I. Fungicides registered for use against diseases of foliage plants in Canada

Plant ¹	Site ²	Disease	Trade Name	Active Ingredient	Formulation	PCP#
Adiantum	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Aglaonema	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Aliette	fosetyl-Al	80% WDG	24458
			Truban	etridiazole	30% WP	11460
			Truban	etridiazole	25% EC	12222
			Subdue Maxx	metalaxyl	240 g/L	27055
	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
Anthurium	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Truban	etridiazole	30% WP	11460
			Truban	etridiazole	25% EC	12222
	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
Aphelandra	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Exotherm Termil	chlorothalornil	20% PO	16661
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
			Subdue Maxx	metalaxyl	240 g/L	27055
Asparagus	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Cactus	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
Caladium	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Exotherm Termil	chlorothalornil	20% PO	16661
			Damping-off (<i>Pythium</i> and <i>Phytophthora</i>)	Subdue Maxx	metalaxyl	240 g/L
	GH and IS ³	Damping –off (<i>Rhizoctonia solani</i>)	Compass	trifloxystrobin	50 WG	27527
Chlorophytum	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Cissus	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
		Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
	GH	Root rot (<i>Fusarium</i>)	Mycostop	<i>Streptomyces griseoviridis</i>	WP	26265
Cordiaum	GH	Gray mold	Decree	fenhexamid	50% WDG	26132

APPENDIX I. Fungicides registered for use against diseases of foliage plants in Canada continued

Plant ¹	Site ²	Disease	Trade Name	Active Ingredient	Formulation	PCP#
Dieffenbachia	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Subdue Maxx	metalaxyl	240 g/L	27055
			Truban	etridiazole	30% WP	11460
			Truban	etridiazole	25% EC	12222
Dieffenbachia	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR	27116
					WP	27115
Dieffenbachia	GH	Root rot (<i>Fusarium</i>)	Mycostop	<i>Streptomyces griseoviridis</i>	WP	26265
Dizygotheca	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
Dracaena	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
Dracaena	GH	Fusarium leaf spot	Rovral	iprodione	50% WDG	24709
Echeveria	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Daconi 12787	chlorothalonil	40% SU	15724
			Daconi 12787	chlorothalonil	90% WDG	24794
Echeveria	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Truban	etridiazole	30% WP	11460
			RootShield	<i>Trichoderma harzianum</i>	GR	27116
Echeveria	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	WP	27115
Epipremnum	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Aliette	fosetyl-Al	80% WDG	24458
			Truban	etridiazole	30% WP	11460
Epipremnum	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	Truban	etridiazole	25% EC	12222
Epipremnum	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR	27116
			RootShield	<i>Trichoderma harzianum</i>	WP	27115
Episcia	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
Episcia	GH	Gray mold	Rovral	iprodione	50% WDG	24709
Euonymus	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
Euonymus	GH	Gray mold	Rovral	iprodione	50% WDG	24709
Euphorbia	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
Euphorbia	GH	Gray mold	Rovral	iprodione	50% WDG	24709
Ficus	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
Ficus	GH	Gray mold	Rovral	iprodione	50% WDG	24709

¹See Appendix II for a list of Latin and common names of foliage plants.

²GH = Greenhouse

³IS = Interiorscape

APPENDIX I. Fungicides registered for use against diseases of foliage plants in Canada continued

Plant ¹	Site ²	Disease	Trade Name	Active Ingredient	Formulation	PCP#
Fittonia	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Gynura	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Hedera		Colletotrichum leaf spot	Dithane M-45 Dithane WSP Manzate 200	mancozeb mancozeb mancozeb	80% WP 80% WP 75% DF	8556 23655 21057
	GH	Damping –off (<i>Pythium</i> and <i>Phytophthora</i>)	Subdue Maxx	metalaxyl	240 g/L	27055
	GH	Gray mold	Decree Rovral Rovral	fenhexamid iprodione iprodione	50% WDG 50% WP 50% WDG	26132 15213 24709
Hypoestes	GH	Damping –off (<i>Rhizoctonia solani</i>)	Compass	trifloxystrobin	50% WG	27527
		Colletotrichum leaf spot	Protect T/O Penncozeb	mancozeb mancozeb	80% WP 80%	25553 25396
		Gray mold	Decree Rovral Rovral	fenhexamid iprodione iprodione	50% WDG 50% WP 50% WDG	26132 15213 24709
Kalanchoe	GH	Gray mold	Decree Phyton Rovral Rovral	Fenhexamid copper iprodione iprodione	50% WDG 5.5% 50% WP 50% WDG	26132 21699 15213 24709
Maranta	GH	Helminthosporium leaf spot	Daconi 12787 Daconi 12787	chlorothalonil chlorothalonil	40% SU 90% WDG	15724 24794
	GH	Gray mold	Decree Rovral Rovral	fenhexamid iprodione iprodione	50% WDG 50% WP 50% WDG	26132 15213 24709
Monstera	GH	Gray mold	Decree Rovral Rovral	fenhexamid iprodione iprodione	50% WDG 50% WP 50% WDG	26132 15213 24709
Nephrolepis	GH	Gray mold	Decree Rovral Rovral	fenhexamid iprodione iprodione	50% WDG 50% WP 50% WDG	26132 15213 24709
Palms	GH and IS	Damping-off (<i>Rhizoctonia solani</i>)	Compass	trifloxystrobin	50% WG	27527
	GH	Fusarium wilt	Mycostop	<i>Streptomyces griseoviridis</i>	WP	26265
	GH	Gray mold	Decree Exotherm Termil	fenhexamid chlorothalonil	50% WDG 20% PO	26132 16661
	GH	Helminthosporium leaf spot	Daconi 12787 Daconi 12787	chlorothalonil chlorothalonil	40% SU 90% WDG	15724 24794
Peperomia	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Subdue Maxx Truban Truban	metalaxyl etrizadiazole etrizadiazole	240 g/L 30% WP 25% EC	27055 11460 12222

Plant¹	Site²	Disease	Trade Name	Active Ingredient	Formulation	PCP#
	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
	GH	Gray mold	Decree Rovral Rovral	fenhexamid iprodione iprodione	50% WDG 50% WP 50% WDG	26132 15213 24709

¹See Appendix II for a list of Latin and common names of foliage plants.

²GH = Greenhouse

³IS = Intiorscape

APPENDIX I. Fungicides registered for use against diseases of foliage plants in Canada continued

Plant ¹	Site ²	Disease	Trade Name	Active Ingredient	Formulation	PCP#
Philodendron	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Subdue Maxx	metalaxyl	240 g/L	27055
			Truban	etridiazole	30% WP	11460
			Truban	etridiazole	25% EC	12222
	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
Philodendron	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Philodendron	GH	Phytophthora blight; Dactylaria leaf spot	Daconil 2787	chlorothalonil	40% SU	15724
			Daconil 2787	chlorothalonil	90% WDG	24794
Pilea	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Truban	etridiazole	30% WP	11460
			Truban	etridiazole	25% EC	12222
			RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
Pilea	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Polyscias	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
Rhoeo	GH	Tan leaf spot	Daconil 2787	chlorothalonil	40% SU	15724
			Daconil 2787	chlorothalonil	90% WDG	24794
Rumohra	GH	Ascochyta blight; Rhizoctonia blight; Cercospora leaf spot; Cylindrocladium leaf spot	Daconil 2787	chlorothalonil	40% SU	15724
			Daconil 2787	chlorothalonil	90% WDG	24794
Saxifraga	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Schefflera	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
			Aliette	fosetyl-Al	80% WDG	24458
			Subdue Maxx	metalaxyl	240 g/L	27055
Senecio	GH	Gray mold	Decree	fenhexamid	50% WDG	26132
			Rovral	iprodione	50% WP	15213
			Rovral	iprodione	50% WDG	24709
Spathiphyllum	GH	Damping-off; root & stem rots (<i>Pythium</i> & <i>Phytophthora</i>)	Aliette	fosetyl-Al	80% WDG	24458
			RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
Spathiphyllum	GH	Damping-off; root rots (<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>)	RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
			RootShield	<i>Trichoderma harzianum</i>	GR WP	27116 27115
Syngonium	GH	Cephalosporium leaf spot	Daconil 2787	chlorothalonil	40% SU	15724
			Daconil 2787	chlorothalonil	90% WDG	24794

Plant ¹	Site ²	Disease	Trade Name	Active Ingredient	Formulation	PCP#
--------------------	-------------------	---------	------------	-------------------	-------------	------

¹See Appendix II for a list of Latin and common names of foliage plants.

²GH = Greenhouse

³IS = Interiorscape

APPENDIX II. Latin and common names of foliage plants

LATIN NAME	COMMON NAME
<i>Aglaonema spp.</i>	Chinese evergreen
<i>Aphelandra squarrosa</i>	Zebra plant
<i>Araucaria heterophylla</i>	Norfolk Island pine
<i>Asparagus densiflorus</i>	Sprengeri asparagus
<i>Brassaia actinophylla</i>	Schefflera; Umbrella tree
Cacti - <i>Opuntia ficus-indica</i>	Spineless cactus
- <i>Opuntia vulgaris</i>	Prickly pear cactus
<i>Caladium spp.</i>	Caladium
<i>Chlorophytum comosum</i>	Spider plant
<i>Cissus antartica</i>	Kangaroo vine
<i>Cissus rhombifolia</i>	Grape ivy
<i>Cordiaum variegatum</i>	Croton
<i>Dieffenbachia spp.</i>	Dieffenbachia
<i>Dizygotheca elegantissima</i>	False aralia
<i>Dracaena deremensis</i>	Janet Craig dracaena
<i>Dracaena deremensis</i>	Warneckii dracaena
<i>Dracaena fragrans</i>	Corn plant
<i>Dracaena fragrans</i>	Massangeana cane
<i>Dracaena marginata</i>	Madagascar dragon tree
<i>Dracaena reflexa</i>	Pleomele
<i>Dracaena sanderana</i>	Ribbon plant
<i>Dracaena surculosa</i>	Gold dust dracaena
<i>Echeveria carnicolor</i>	Coral echeveria
<i>Epipremnum aureum</i>	Pothos; Devil's ivy
<i>Episcia spp.</i>	Flame violets
<i>Euonymus spp.</i>	Enonymous
<i>Euphorbia millii</i>	Crown-of-thorns
<i>Fatsia japonica</i>	Japanese aralia
Ferns - <i>Adiantum spp.</i>	Maidenhair fern
- <i>Asplenium nidus</i>	Bird's-nest fern
- <i>Nephrolepis exaltata</i>	Boston fern; Sword fern
- <i>Rumohra adiantiformis</i>	Leatherleaf fern
<i>Ficus benjamina</i>	Weeping fig
<i>Ficus elastica</i>	India rubber tree
<i>Ficus lyrata</i>	Fiddle-leaf fig
<i>Ficus pumila</i>	Creeping fig
<i>Ficus retusa</i>	Cuban laurel fig
<i>Fittonia verschaffetii</i>	Silver-nerve plant

APPENDIX II. (continued)

LATIN NAME	COMMON NAME
<i>Gynura aurantiaca</i>	Purple passion
<i>Hedera helix</i>	English ivy
<i>Hypoestes spp.</i>	Pink splash
<i>Kalanchoe tomentosa</i>	Pussy ears; Panda plant
<i>Maranta leuconeura</i>	Prayer plant
<i>Monstera deliciosa</i>	Cutleaf philodendron
<i>Nephtytis</i> (see <i>Syngonium</i>)	
Palms - <i>Caryota mitis</i>	Fishtail palm
- <i>Chamaedorea elegans</i>	Parlor palm; Neanthe bella
- <i>Chamaedorea erumpens</i>	Bamboo palm
- <i>Chamaedorea seifrizii</i>	Reed palm
- <i>Chamaedorea humilis</i>	European fan palm
- <i>Chrysalidocarpus lutescens</i>	Areca palm
- <i>Cycas revoluta</i>	Sago palm
- <i>Howea forsterana</i>	Kentia palm; Sentry palm
- <i>Livistona chinensis</i>	Chinese fan palm
- <i>Phoenix roebelini</i>	Pygmy date palm
- <i>Ptychosperma elegans</i>	Solitaire palm; Alexander palm
- <i>Rhapis excelsa</i>	Lady palm
- <i>Rhapis humilis</i>	Slender lady palm
<i>Peperomia spp.</i>	Peperomia
<i>Philodendron scandens oxycardium</i>	Heart-leaf philodendron
<i>Philodendron selloum</i>	Lacy-tree philodendron
<i>Pilea cadierei</i>	Aluminium plant
<i>Pilea microphylla</i>	Artillary plant
<i>Polyscias spp.</i>	Aralia
<i>Rhoeo spatheca</i>	Moses-in-the-cradle
<i>Rumohra adiantiformis</i>	Leatherleaf fern
<i>Sansevieria spp.</i>	Sword plant; Sansevieria
<i>Saxifraga spp.</i>	Rockfoil; Saxifrage
<i>Scheffera actinophylla</i>	(see <i>Brassaia actinophylla</i>)
<i>Scheffera arbuticola</i>	Hawaiian schefflera; Dwarf schefflera
<i>Scindapsus aureus</i>	(see <i>Epipremnum aureum</i>)
<i>Senecio spp.</i>	Cineraria
<i>Spathiphyllum spp.</i>	Peace lily
<i>Syngonium spp.</i>	Nephtytis; Arrowhead vine
<i>Yucca spp.</i>	Yucca